

# Impact of removal of invasive species *Kappaphycus alvarezii* from coral reef ecosystem in Gulf of Mannar, India

B. Kamalakannan<sup>1</sup>, J. Joyson Joe Jeevamani<sup>1</sup>, N. Arun Nagendran<sup>2</sup>,  
D. Pandiaraja<sup>2</sup> and S. Chandrasekaran<sup>3,\*</sup>

<sup>1</sup>Department of Botany, and <sup>2</sup>National Centre of Excellence in Statistical and Mathematical Modelling in Bioresources Management (MHRD), Thiagarajar College, Madurai 625 009, India

<sup>3</sup>Department of Plant Sciences, School of Biological Sciences, Madurai Kamaraj University, Madurai 625 021, India

***Kappaphycus alvarezii* is a commercially important red alga being intentionally introduced in marine waters worldwide for the production of kappa carrageenan. Its introduction into the Gulf of Mannar Biosphere Reserve during the 1990s and its subsequent escape from cultivation sites have paved the way for its invasion into the coral reef ecosystem of Kurusadai Island. Since the report of its invasion in 2008, removal of *K. alvarezii* from the reefs has been started by means of manual removal (hand plucking). This article details the unsuccessful attempt and negative impact of the eradication programme. Regrowth of *K. alvarezii* from removal points and drifting broken fragments resulting during removal have led to further establishment in the reef environment. Variation in the morphology of *K. alvarezii* populations after their removal has been observed. A significant reduction in the cover of coral and native algae due to the increase in abundance of *K. alvarezii* was evident from the study. The need for immediate scientific control measures to eradicate the invasive alga is discussed.**

**Keywords:** Coral reefs, Gulf of Mannar, invasive species, *Kappaphycus alvarezii*, manual removal.

BIODIVERSITY is affected by the invasion of exotic species in new geographical locations. The introduction and spread of non-native species have significantly altered the ecological functions of marine ecosystems<sup>1,2</sup>. Exotic marine algae which behave as invasive species have impacted the native coral communities at the sites of incursion<sup>3-5</sup>. One such alga is *Kappaphycus alvarezii* (Doty) Doty ex. P. Silva (Rhodophyta: Solieriaceae). It is one of many seaweeds being intentionally introduced for the production of kappa carrageenan worldwide<sup>6</sup>. The farming of *K. alvarezii* was initiated in the Philippines during 1960s with local varieties of its wild populations, and it has expanded further to other parts of the world with different cultivation technologies<sup>7,8</sup>. However,

*K. alvarezii* poses serious threats to native corals through overgrowing and smothering<sup>9,10</sup>.

The commercial cultivation of *K. alvarezii* in India was strongly opposed due to the prediction of its likely invasiveness<sup>11</sup>, as it is exotic to Indian marine environments. However, field vigilance and environmental impact assessments showed no visible harmful effects from this alga<sup>12</sup> and thus its cultivation continues. However, later studies had shown its smothering effect on live corals in Kurusadai Island in the Gulf of Mannar (GoM)<sup>13,14</sup>. Incidents of *K. alvarezii* invasion on corals in GoM<sup>13</sup> were reported<sup>15,16</sup>. There were immediate remedial responses from the State Government organizations to control/eradicate the alga.

Control of *K. alvarezii* in invaded communities has been carried out by either physical or biological methods<sup>17</sup>, or both. An underwater vacuuming system, Super Sucker, has been used as a physical method to eradicate *K. alvarezii* in Kane'ohe Bay, O'ahu. Also, native collector urchins (*Tripneustes gratilla*) were used to control *K. alvarezii*, which clear them through grazing. However, in Kurusadai Island, the Forest Department decided to use the manual removal method to reduce the impact of *K. alvarezii* on corals. In this article, we report on the consequences of the *K. alvarezii* removal process carried out in GoM.

## Materials and methods

### Study area

Kurusadai Island (9°15'N; 79°12'E) is a part of Mandapam group islands in GoM biosphere reserve, Tamil Nadu (for a map of the study area see ref. 13). Study sites are part of continuous fringing reefs located on the southern side of the Island which extend up to 500 m with varying depths of 0.5–2.0 m. A survey in 2005 revealed the presence of 54.9% live coral cover in the Island<sup>18</sup>. The coral ecosystem has experienced a recent coral–algal phase shift<sup>19</sup> due to coral bleaching in 2010.

\*For correspondence. (e-mail: krishnanasc@yahoo.co.in)

The present study has continued from 2007 after the preliminary report of *K. alvarezii* invasion on patches of corals in intertidal region of the Island<sup>13</sup>. To examine the effects of *K. alvarezii* invasion on native coral reef communities, we carried out subsequent monitoring with the addition of control ecosystems devoid of *K. alvarezii* invasion. The control ecosystems are situated west of the invaded ecosystems and they are separated by a distance of about 750–1000 m. The invaded and control ecosystems were further divided into two sites, viz. site 1 (50 m from the shore) and site 2 (100 m from the shore). Depth at site 1 varies from 0.5 to 1.0 m and that at site 2 varies from 1.0 to 2.0 m for both ecosystems. Other than depth, all other biotic and abiotic factors are common to both sites of control and invaded ecosystems, except the presence of *K. alvarezii* in the invaded ecosystem.

### Benthic community analysis

Estimates of the coral cover, live cover of *K. alvarezii* on corals, live cover of native algae and sand/rubble (expressed in %) were based on 80 randomly placed, 1 m<sup>2</sup> quadrats ( $n = 20$  per site per ecosystem). Stratified sampling was adopted to select *K. alvarezii*-invaded coral colonies in the study sites (sites 1 and 2) of the invaded ecosystems. Changes in abundance of *K. alvarezii* and native algae were estimated independently using 1 m<sup>2</sup> quadrats without segregation of sites ( $n = 20$  per ecosystem). Quadrats were located with a GPS (Garmin, Taiwan) and were visited periodically once in a month (between March and August) during 2008–2012. All quadrats were at least 1 m apart from each other. Parameters such as species abundance, species richness and evenness were estimated to study algal dynamics. Species abundance was calculated as the ratio between total number of individuals of a species and the number of quadrats in which the species were present. Species richness ( $S$ ) was determined for each quadrat as the number of identified algal taxa. Simpson index of diversity ( $1 - D$ ), which measures the probability of any two individuals randomly drawn from a community belonging to the same species, was used as a measure of species richness and evenness. For each quadrat, it was calculated as

$$1 - D = 1 - \sum_{i=1}^S [n_i(n_i - 1) / N(N - 1)],$$

where  $S$  is the species richness,  $n_i$  is the number of individuals in the  $i$ th species, and  $N$  is the total number of individuals of all species present in a quadrat. The value of  $1 - D$  ranges from 0 to 1 and thus a higher value indicates greater diversity.

Though manual removal of *K. alvarezii* was started from 2009 (discussed below), all quadrats used for benthic community analysis in the invaded ecosystems

were not affected by the manual removal process during 2009–2012. Due to mass eradication in 2012, these quadrats were not monitored further.

### Assessment of manual removal impact on *K. alvarezii*

In early March 2009, the Tamil Nadu Forest Department initiated the removal of *K. alvarezii* by manual method (hand plucking) from invaded coral colonies at intertidal zone in Kurusadai Island. We did the first survey on the same day (8 March 2009). By visual survey, we selected nine removal points, i.e. reef substrates from which the alga was completely removed. Each removal point was covered by a GPS-marked quadrat (1 m<sup>2</sup>) to analyse the *K. alvarezii* regrowth pattern. Prior to marking, we estimated area (cm<sup>2</sup>/m<sup>2</sup>) and well-drained fresh biomass (kg/m<sup>2</sup>) of removed *K. alvarezii* colonies from the respective quadrats, which were noted as initial measurements. Resurvey was done at marked quadrats in early September 2009 (1 September 2009) spanning an time-interval of 175 days and estimated values were taken as final measurements. Algal colonies were removed during low-tide condition following the method of Conklin and Smith<sup>20</sup>. Daily growth rate (DGR) was estimated from the collected biomass using the equation given by Rueness *et al.*<sup>21</sup>

$$\text{Growth rate (\% day}^{-1}\text{)} = 100 \ln (W_t/W_0)/t,$$

where  $W_0$  is initial weight,  $W_t$  final weight;  $t$  is the time-interval (days).

Qualitative data were taken using photographs. All observations and estimations were done during low-tide conditions.

### Statistical analysis

Percentage cover data did not satisfy the assumptions of normal distribution, tested using Shapiro–Wilk’s test, even after transformations. Mann–Whitney  $U$ -test was applied to test the null hypothesis stating that the percentage cover of each benthic component in control and invaded ecosystems, treating them as independent samples, has the same median. Since the data are non-normal, we used the median measure instead of mean for comparisons. Wilcoxon signed ranks test was used to test whether the median of each benthic variable differs significantly between successive years since 2008. However, colony area and biomass data of *K. alvarezii* collected for manual removal assessment were normally distributed. Hence, we used paired samples  $t$ -test to reveal differences in their respective means between pre- and post-removal period. All statistical analyses were performed with IBM SPSS Statistics version 20.0.0.

## Results

### Benthic community analysis

We found no significant difference in the median of the studied benthic variables between successive years in both sites of control ecosystem (Wilcoxon signed ranks test:  $P > 0.05$ ,  $N = 20$ , in all cases; Table 1). Compared to the control ecosystems, a remarkable decline in coral cover was observed in the *K. alvarezii*-invaded ecosystems ( $U = 4276.0$ ,  $P < 0.01$ ; Figure 1). In *K. alvarezii*-invaded ecosystems, median of coral cover was reduced from 64.9% in 2008 to 33.4% in 2012 and 45.6% in 2008 to 0% in 2012 at sites 1 and 2 respectively (Figure 1). A reduction in median of coral cover was significant between successive years at both sites of *K. alvarezii*-invaded ecosystems. *K. alvarezii* showed an increasing trend in its distribution at both sites of invaded ecosystem (Figure 1). Among the sites of the invaded ecosystem, median of *K. alvarezii* cover on corals had attained the maximum (97.6%) in site 2 rather than site 1 (60.9%) during 2012. It also showed significant difference in the median of *K. alvarezii* cover on invaded corals during successive years at both sites (Table 1).

During the study period, native algal species also experienced significant reduction in their cover at the *K. alvarezii*-invaded ecosystems compared to the control ecosystems ( $U = 9009.0$ ,  $P < 0.01$ ). Significant reduction in median of native algal cover was observed from the year 2011 in site 1 and from 2010 in site 2 of *K. alvarezii*-invaded ecosystem (Table 1). The median number of algal species ( $S$ ) per quadrat (*K. alvarezii* not included) was 7 (range: 3–11 species,  $n = 100$ ) in the control ecosystems and 5 (range: 1–8 species) in the invaded ecosys-

tems. Median of species richness ( $S$ ) showed significant difference between control and invaded ecosystems ( $U = 2031.5$ ,  $P < 0.01$ ). In the invaded ecosystem, drastic decline in species richness ( $S$ ) was observed during post-removal period (Table 2). Simpson diversity ( $1 - D$ ) per quadrat (without *K. alvarezii*) was found to be significantly higher ( $U = 3457.0$ ,  $P < 0.01$ ) in control ecosystems (median = 0.8,  $n = 100$ ) than invaded ecosystems (median = 0.7,  $n = 100$ ). Further, Simpson diversity in *K. alvarezii*-invaded quadrats reduced gradually during post-removal period (Table 2). This reduction clearly depicted the increase in dominance of *K. alvarezii* in the invaded ecosystems. Due to increase in abundance, *K. alvarezii* had shifted from lower rank (9th) during pre-removal to higher rank (3rd) and subsequently attained top rank (1st) after 2009, i.e. post-removal. Consequently, dominant native species such as *Gracilaria* sp., *Gelidiella* sp., *Caulerpa* sp. and *Padina* sp. were not recorded from the *K. alvarezii*-invaded ecosystems and other species such as *Sargassum* sp., *Turbinaria* sp., *Halimeda* sp., *Ulva reticulata* and *Hypnea* sp. had declined considerably (Table 2).


### Manual removal impacts on *K. alvarezii*

In the invaded ecosystem, the estimated initial (pre-removal) mean colony area of *K. alvarezii* was  $1457.44 \pm 113.11$  cm<sup>2</sup>/m<sup>2</sup> (mean  $\pm$  SE,  $n = 9$ ). Its biomass was estimated to be  $4.52 \pm 0.32$  kg/m<sup>2</sup>. We observed extensive regrowth of *K. alvarezii* at the removal points (Figure 2 a). After 175 days, the mean colony area and biomass of *K. alvarezii* were found to be  $3382.67 \pm 192.09$  cm<sup>2</sup>/m<sup>2</sup> and  $12.77 \pm 0.68$  kg/m<sup>2</sup> respectively. Paired *t*-test revealed significant difference between pre- and post-removal

**Table 1.** Comparison of benthic cover components between successive years in control and *K. alvarezii*-invaded ecosystems of Kurusadai Island, Gulf of Mannar (GoM) during 2008–2012

Ecosystem	Site	Benthic cover	Comparison between successive years ( $P$ value)			
			2008 vs 2009	2009 vs 2010	2010 vs 2011	2011 vs 2012
Control	Site 1	Coral	0.520	0.243	0.601	0.654
		<i>K. alvarezii</i> on coral	–	–	–	–
		Native algae	0.845	0.248	0.627	0.136
	Site 2	Sand/rubble	0.910	0.913	0.825	0.334
		Coral	0.918	0.165	0.856	0.526
		<i>K. alvarezii</i> on coral	–	–	–	–
Invaded	Site 1	Native algae	0.756	0.070	0.376	0.433
		Sand/rubble	0.609	0.983	0.446	0.765
		Coral	<b>0.010</b>	<b>0.005</b>	<b>0.013</b>	<b>0.010</b>
	Site 2	<i>K. alvarezii</i> on coral	<b>0.001</b>	<b>0.000</b>	<b>0.000</b>	<b>0.000</b>
		Native algae	0.388	0.062	<b>0.000</b>	<b>0.000</b>
		Sand/rubble	0.209	0.051	<b>0.008</b>	<b>0.028</b>
		Coral	<b>0.002</b>	<b>0.001</b>	<b>0.002</b>	<b>0.031</b>
		<i>K. alvarezii</i> on coral	<b>0.001</b>	<b>0.001</b>	<b>0.002</b>	<b>0.006</b>
		Native algae	0.814	<b>0.001</b>	<b>0.001</b>	<b>0.005</b>
	Site 2	Sand/rubble	0.109	0.109	0.109	0.273

Bold values show significant difference at  $\alpha = 0.05$  level.  $P$  values are based on Wilcoxon signed ranks test.



**Figure 1.** Distribution (% cover) of different benthic components in site 1 (50 m from the shore) and site 2 (100 m from the shore) of both control and *K. alvarezii*-invaded ecosystems in Kurusadai Island, Gulf of Mannar (GoM) during 2008–2012. Box plots show the minimum, 25th percentile, median, 75th percentile and maximum values (outliers not shown). White box represents control ecosystems and filled box represents invaded ecosystems.

groups of *K. alvarezii* for colony area ( $t = 23.5$ ,  $P < 0.01$ ) and biomass ( $t = 22.1$ ,  $P < 0.01$ ). After removal, an increase of ca. 132% and 182% was recorded in colony area and biomass of *K. alvarezii* respectively. Estimated DGR of *K. alvarezii* for 175 days was 0.6%. Colonies of *K. alvarezii* fragmented from manual eradication attached themselves to neighbouring healthy coral colonies, and then expanded further (Figure 2b). Wave action was also responsible for the reintroduction of removed and discarded *K. alvarezii* colonies from the seashore to uninvaded corals (Figure 2c). Washed-up fragments of *K. alvarezii* colonies were observed along the seashore (Figure 2d). Twelve algal clumps with varying biomass of 150–1125 g

(fresh weight) were collected along the shore of the invaded ecosystem during the study period. In addition to these observations and estimations, we observed a novel adaptation of the species after the interference of manual removal. In 2008, the growth of *K. alvarezii* was reported to be as a green mat over the top and lateral sides of the corals (Figure 2e) and the major axis of *K. alvarezii* closely adhered with the rough surface of the corals in the study region<sup>13</sup>. In contrast, here we noticed unusual dome-like growth of *K. alvarezii* along the coral landscape (Figure 2f), particularly on manually damaged *K. alvarezii* colonies. The dome protruded out of the coral landscape and its height was  $10 \pm 0.76$  cm in site 1 and

**Table 2.** Species abundance, richness, diversity and evenness of identified algal taxa in control and *K. alvarezii*-invaded ecosystems of Kurusadai Island, GoM during 2008–2012

Algal species	Control ecosystem					Invaded ecosystem				
	Pre-removal		Post-removal			Pre-removal		Post-removal		
	2008	2009	2010	2011	2012	2008	2009	2010	2011	2012
Species abundance <sup>a</sup>										
<i>Sargassum</i> sp.	14.9 (4)	18.9 (3)	23.9 (1)	15.3 (2)	19.0 (1)	11.9 (1)	10.6 (1)	4.7 (2)	2.4 (4)	1.8 (2)
<i>Gracilaria</i> sp.	21.8 (2)	22.4 (1)	23.7 (2)	15.3 (2)	16.2 (2)	11.2 (2)	5.9 (2)	5.5 (1)	1.0 (8)	0 (6)
<i>Gelidella</i> sp.	22.4 (1)	14.8 (4)	12.8 (4)	14.5 (4)	9.3 (5)	8.6 (3)	5.3 (4)	1.4 (8)	0 (10)	0 (6)
<i>Turbinaria</i> sp.	17.4 (3)	20.9 (2)	20.9 (3)	19.1 (1)	13.6 (3)	6.3 (4)	5.2 (5)	2.5 (5)	1.2 (7)	1.0 (4)
<i>Caulerpa racemosa</i>	8.0 (6)	3.9 (7)	5.5 (5)	9.5 (5)	5.8 (7)	4.6 (5)	3.5 (7)	2.9 (4)	2.6 (3)	0 (6)
<i>Caulerpa taxifolia</i>	9.5 (5)	2.3 (9)	2.5 (8)	4.0 (8)	5.9 (6)	4.5 (6)	5.4 (3)	0 (10)	2.0 (5)	0 (6)
<i>Padina</i> sp.	7.0 (8)	3.7 (8)	4.6 (6)	8.9 (6)	10.9 (4)	4.3 (7)	1.3 (10)	1.0 (9)	1.0 (8)	0 (6)
<i>Halimeda</i> sp.	2.6 (10)	1.0 (10)	1.5 (10)	1.2 (11)	1.9 (9)	0 (10)	3.6 (6)	1.9 (6)	0 (10)	1.0 (4)
<i>Ulva reticulata</i>	8.0 (6)	4.8 (5)	3.7 (7)	4.6 (7)	5.3 (8)	3.1 (8)	3.0 (8)	1.7 (7)	4.1 (2)	1.4 (3)
<i>Hypnea</i> sp.	7.0 (8)	4.2 (6)	1.4 (11)	1.7 (9)	1.0 (11)	0 (10)	0 (11)	0 (10)	1.3 (6)	0 (6)
<i>Dictyosphaeria cavernosa</i>	1.1 (11)	1.0 (10)	1.6 (9)	1.3 (10)	1.4 (10)	0 (10)	0 (11)	0 (10)	0 (10)	0 (6)
<i>Kappaphycus alvarezii</i>	–	–	–	–	–	2.1 (9)	2.5 (9)	3.5 (3)	4.2 (1)	4.7 (1)
Species richness ( <i>S</i> ) <sup>b</sup>	5.5 (2.7)	6.0 (2.7)	7.0 (2.0)	7.0 (1.0)	8.0 (1.0)	6.0 (1.7)	5.5 (1.7)	4.5 (2.0)	4.0 (2.0)	2.0 (1.7)
Simpson index of diversity ( $1 - D$ ) <sup>b</sup>	0.8 (0.1)	0.7 (0.1)	0.8 (0.1)	0.8 (0.04)	0.8 (0.05)	0.8 (0.1)	0.8 (0.1)	0.7 (0.2)	0.5 (0.6)	0.4 (0.9)

<sup>a</sup>Values denote abundance (species rank); <sup>b</sup>Values represent median (interquartile range;  $n = 20$  quadrats for each year); *S* and  $1 - D$  values of invaded ecosystem devoid of *K. alvarezii* data.

18.6 ± 1.56 cm in site 2 ( $n = 10$  per site) of the invaded ecosystem.


During our recent field visit in early March 2013, there appeared a widespread occurrence of *K. alvarezii* at both study sites of the invaded ecosystem which can be viewed clearly at low tides (Figure 3a). The number of *K. alvarezii* colonies was found to be 8.5 and 11.2 (per m<sup>2</sup>;  $n = 10$  quadrats) in sites 1 and 2 of invaded ecosystem respectively. It is significantly higher than the average number of *K. alvarezii* colonies observed in the corresponding sites of invaded ecosystem during pre-removal, i.e. 2009 (Table 3). In 2013, estimates of *K. alvarezii* colony area had ranged from 9.0 to 1716.0 cm<sup>2</sup> (mean = 502.7 cm<sup>2</sup>,  $n = 85$  colonies) in site 1 and 1.0 to 1722.0 cm<sup>2</sup> (mean = 239.3 cm<sup>2</sup>;  $n = 112$  colonies) in site 2 of the invaded ecosystems. Compared to mean *K. alvarezii* colony area during pre-removal, a significant reduction in mean colony area was observed in both sites of invaded ecosystems during post-removal, i.e. 2013 (Table 3; Figure 3b).

## Discussion

The results showed that innate benthos (such as corals, native algae) of Kurusadai Island were not altered much between sites as well as removal periods in the control ecosystems. A minimum level of reduction in coral cover of control ecosystems was observed during the study period (Figure 1). Competition between corals and native algae could be a reason for such reduction. However, such effects are common and fluctuate due to interannual

variations. Absence of the invasive species *K. alvarezii* in the control ecosystems could be a specific reason for such intact ecosystem, whereas innate benthos cover was considerably decreased due to increase in the cover and abundance of *K. alvarezii* in the invaded ecosystems. Failure to restore pre-invasion status and predominant increase in *K. alvarezii* cover at invaded ecosystems could suggest that the removal has been counter-productive. Vigorous regrowth and establishment of *K. alvarezii* population has been observed after the removal process.

The reason for manual removal failure may be due to (i) biology of the invasive species and (ii) inefficacy of the method employed to eradicate *K. alvarezii*. The alga affects the corals by completely smothering them<sup>13,14</sup>. Consequently, there could be every possibility for leaving algal fragments within the coral tissues during removal. We observed small fragments of *K. alvarezii* firmly attached to the corals (Figure 3c), evidently detached during removal. The alga has the ability to coalesce into the tissues of corals, which provides a strong means of attachment<sup>9</sup>. Thus pieces of algal remains inside the coral tissue could facilitate regrowth under ambient natural conditions. Change in the structural form of *K. alvarezii* (i.e. dome protuberance) was observed only after manual removal. This modification may be due to non-availability of the substrate (i.e. corals) near the replenishing *K. alvarezii*-invaded coral colonies. As a result, such algal colonies have adapted themselves to extend their colony expansion vertically. Most likely such adaptation still favours its spread by break and drift of protruding branches as fragments due to wave action. Even the smallest




**Figure 2.** *a*, Extensive regrowth of *K. alvarezii* from a removal point; *b*, Reattachment of drifted algal fragments to live corals; *c*, Rebound of a discarded massive algal colony into reef substrate through wave action; *d*, Washed-up algal fragments found along the seashore; *e*, *K. alvarezii* smothering corals by growing as elastic rubber sheet (from Chandrasekaran *et al.*<sup>15</sup>); *f*, Transition from sheet like appearance to dome-like structure, showing its extraordinary phenotypic plasticity.

fragments of size 0.05 g will be able to disperse and establish themselves widely if given sufficient time<sup>9</sup>.

The method employed for this alga removal (i.e. hand plucking) was not well planned and was unscientific. Available field experimental data on manual removal of *K. alvarezii* from corals in Hawaii<sup>20</sup> have shown that the process is a daunting task. In the Gulf of Mannar National Park, the park managers have attempted eradication operations and they removed *K. alvarezii* attached to coral colonies in a ‘pluck and throw’ manner with no proper strategy to dispose or prevent spread of broken fragments. We observed several *Acropora* coral fragments attached to *K. alvarezii* both on-site (reef substrates) and off-site (seashore; Figure 3 *d* and *e*). To date, local island managers are collecting the washed-up coral

fragments with *K. alvarezii* and dumping them on the seashore. So far, it is considered that the species could spread or reproduce by vegetative fragmentation because sexual reproduction through spores is rare and not viable<sup>22,23</sup>. Hence fragmentation resulting from manual removal facilitates dispersal of fragments, and such broken fragments seem to attach and re-establish in the reef ecosystem. Widespread observation of higher number as well as small-sized colonies of *K. alvarezii* in invaded ecosystems during our recent visit in 2013 has supported this claim (Figure 3 *b*). Strong regeneration ability of this species and unscientific eradication strategies have contributed to significant spread of invasion in the locality. Similar negative feedback has been accounted in physical removal of invasive alga *Sargassum muticum* in England<sup>24</sup>.



**Figure 3.** *a*, Photographic view showing *Kappaphycus* algal mounds over corals in March 2013; *b*, Numerous small-sized colonies of *K. alvarezii* resulted from manual eradication; *c*, Small-sized algal piece (1 cm) reattached to corals after removal; *d*, *e*, Drifted coral fragments found on reefs and seashore.

**Table 3.** Frequency and colony size ( $\text{cm}^2/\text{m}^2$ ) of *K. alvarezii* in invaded ecosystems of Kurusadai Island, GoM during the years 2009 and 2013

Parameter	<i>K. alvarezii</i> invaded ecosystem	Pre-removal (2009)	Post-removal (2013)	<i>t</i> -test**	<i>P</i> value##
No. of colonies*	Site 1	$1.8 \pm 0.4$	$8.5 \pm 2.0$	14.53	< 0.05
	Site 2	$2.2 \pm 0.5$	$11.2 \pm 5.0$	8.02	< 0.05
Area of colonies#	Site 1	$2238.0 \pm 1268.5$	$502.7 \pm 389.0$	10.77	< 0.05
	Site 2	$5855.0 \pm 3102.2$	$239.3 \pm 349.9$	18.82	< 0.05

Data represent mean  $\pm$  SD; \*Number per  $1 \text{ m}^2$  quadrat ( $n = 20$  for 2009;  $n = 10$  for 2013); #  $n = 20$  colonies for site 1 and site 2 in 2009;  $n = 85$  colonies for site 1 and  $n = 112$  colonies for site 2 in 2013; \*\*Unequal sample size *t*-test; ## Significance tested at  $\alpha = 0.05$  level.

Invasive species as ‘ecosystem modifiers’ modify, create and maintain new physico-chemical conditions for their comfortable growth and continued expansion<sup>25</sup>. Their removal should be viewed in the whole ecosystem or community context<sup>26,27</sup>. Hence unscientific control strategies would exacerbate existing complexities and issues.

### Conclusion and implication

This study illustrates the negative effects of the ongoing unscientific manual removal of *K. alvarezii* from invaded coral reef ecosystem in the Gulf of Mannar Marine National Park. If the alga is not controlled properly, there could be chance for it to invade the control ecosystem which is physically separated by considerable distance

from invaded ecosystem and shares similar biogeographic conditions. To eradicate *K. alvarezii* from Kurusadai Island, it is necessary to develop ecologically viable control measures such as use of native herbivores, and mechanical removal by means of sucking pumps which are non-destructive to native communities, especially corals. While it is almost impossible to eliminate established alien species in marine habitats<sup>28</sup>, it is obvious that prevention would be the option to avoid incidents of invasion.

1. Grosholz, E., Ecological and evolutionary consequences of coastal invasions. *Trends Ecol. Evol.*, 2002, **17**, 22–27.
2. Occhipinti-Ambrogi, A. and Savini, D., Biological invasions as a component of global change in stressed marine ecosystems. *Mar. Pollut. Bull.*, 2003, **46**, 542–551.

3. Piazzì, L., Ceccherelli, G. and Cinelli, F., Threat to macroalgal diversity: effects of the introduced green alga *Caulerpa racemosa* in the Mediterranean. *Mar. Ecol. Prog. Ser.*, 2001, **210**, 149–159.
4. Casas, G., Scrosati, R. and Piriz, M. L., The invasive kelp *Undaria pinnatifida* (Phaeophyceae, Laminariales) reduces native seaweed diversity in Nuevo Gulf (Patagonia, Argentina). *Biol. Invasions*, 2004, **6**, 411–416.
5. Britton-Simmons, K. H., Direct and indirect effects of the introduced alga *Sargassum muticum* on benthic, subtidal communities of Washington State, USA. *Mar. Ecol. Prog. Ser.*, 2004, **277**, 61–78.
6. Bixler, H. J., Recent developments in manufacturing and marketing carrageenan. *Hydrobiologia*, 1996, **326/327**, 35–57.
7. Sulu, R., Kumar, L., Hay, C. and Pickering T., *Kappaphycus* seaweed in the Pacific: review of introductions and field testing proposed quarantine protocols. Secretariat of the Pacific Community, Noumea, 2004, pp. 1–84.
8. Bindu, M. S. and Levine, I. A., The commercial red seaweed *Kappaphycus alvarezii* – an overview on farming and environment. *J. Appl. Phycol.*, 2011, **23**, 789–796.
9. Woo, M., Ecological impacts and interactions of the introduced red alga *Kappaphycus striatum* in Kaneohe Bay, Oahu. Master thesis, Department of Botany, University of Hawai'i, Honolulu, 2000.
10. Smith, J. E., 153 Invasive macroalgae on tropical reefs: impacts, interactions, mechanisms and management. *J. Phycol.*, 2003, **39**, 53.
11. Pereira, N. and Verlecar, X. N., Is Gulf of Mannar heading for marine bioinvasion? *Curr. Sci.*, 2005, **89**, 1309–1310.
12. Tewari, A. *et al.*, Environmental impact assessment of *Kappaphycus* cultivation in India in context to global scenario. In *Recent Advances on Applied Aspects of Indian Marine Algae with Reference to Global Scenario* (ed. Tewari, A.), CSMCRI, Bhavnagar, 2006, vol. I, pp. 262–287.
13. Chandrasekaran, S., Nagendran, N. A., Pandiaraja, D., Krishnankutty, N. and Kamalakannan B., Bioinvasion of *Kappaphycus alvarezii* on corals in the Gulf of Mannar, India. *Curr. Sci.*, 2008, **94**, 1167–1172.
14. Kamalakannan, B., Jeevamani, J. J. J., Nagendran, N. A., Pandiaraja, D., Krishnan Kutty, N. and Chandrasekaran, S., *Turbinaria* sp. as victims to *Kappaphycus alvarezii* in reefs of Gulf of Mannar, India. *Coral Reefs*, 2010, **29**, 1077.
15. Anon., Indian coral islands under threat from algae. *Nature*, 2008, **453**, 710–711.
16. Bagla, P., Seaweed invader elicits angst in India. *Science*, 2008, **320**, 1271.
17. <http://dlnr.hawaii.gov/hisc/files/2013/03/Hunter-HISC-final-report-10-07.pdf> (accessed on 4 April 2014).
18. Sukumaran, S., George, R. M. and Kasinathan, C., Biodiversity and community structure of coral reefs around Krusadai Island, Gulf of Mannar, India. *Indian J. Fish.*, 2007, **54**, 275–282.
19. Jeevamani, J. J. J., Kamalakannan, B., Nagendran, N. A. and Chandrasekaran, S., Climate change induced coral bleaching and algal phase shift in reefs of the Gulf of Mannar, India. In *Knowledge Systems of Societies for Adaptation and Mitigation of Impacts of Climate Change* (eds Nautiyal, S. *et al.*), Springer-Verlag, Berlin, 2013, pp. 87–94.
20. Conklin, E. J. and Smith, J. E., Abundance and spread of the invasive red algae, *Kappaphycus* spp., in Kane'ohe Bay, Hawai'i and an experimental assessment of management options. *Biol. Invasions*, 2005, **7**, 1029–1039.
21. Rueness, J., Mathisen, H. A. and Tananger, T., Culture and field observations on *Gracilaria verrucosa* (Huds.) Papenf. (Rhodophyta) from Norway. *Bot. Mar.*, 1987, **30**, 267–276.
22. Zuccarello, G. C., Critchley, A. T., Smith, J., Sieber, V., Lhonneur, G. B. and West, J. A., Systematics and genetic variation in commercial *Kappaphycus* and *Eucheuma* (Solieriaceae, Rhodophyta). *J. Appl. Phycol.*, 2006, **18**, 643–651.
23. Conklin, K. Y., Kurihara, A. and Sherwood, A. R., A molecular method for identification of the morphologically plastic invasive algal genera *Eucheuma* and *Kappaphycus* (Rhodophyta, Gigartinales) in Hawaii. *J. Appl. Phycol.*, 2009, **21**, 691–699.
24. Fletcher, R. L. and Fletcher, S. M., Studies on the recently introduced brown alga *Sargassum muticum* (Yendo) Fensholt. II. Regenerative ability. *Bot. Mar.*, 1975, **18**, 157–162.
25. Crooks, J. A., Characterizing ecosystem-level consequences of biological invasions: the role of ecosystem engineers. *Oikos*, 2002, **97**, 153–166.
26. Zavaleta, E. S., Hobbs, R. J. and Mooney, H. A., Viewing invasive species removal in a whole-ecosystem context. *Trends Ecol. Evol.*, 2001, **16**, 454–459.
27. Hacker, S. D. and Dethier, M. N., Differing consequences of removing ecosystem-modifying invaders: significance of impact and community context to restoration potential. In *Biological Invasions in Marine Ecosystems: Ecological, Management and Geographic Perspectives* (eds Rilov, G. and Crooks, J. A.), Springer-Verlag, Berlin, 2009, pp. 375–385.
28. Thresher, R. E. and Kuris, A. M., Options for managing invasive marine species. *Biol. Invasions*, 2004, **6**, 295–300.

ACKNOWLEDGEMENTS. We thank the Council of Scientific and Industrial Research, New Delhi for financial assistance and Tamil Nadu Forest Department for permission to carry out this work. We thank the two anonymous reviewers for their critical comments and valuable suggestions on the manuscript. We also thank Prof. R. Kannan, Madurai Kamaraj University, Madurai and Mr N. Krishnankutty, Madurai for help during the preparation of the manuscript.

Received 3 October 2013; revised accepted 8 April 2014